



# The Orange Carotenoid Protein: Mechanism of a Photoswitch

Corie Ralston, Sayan Gupta (MBIB)

Cheryl Kerfeld, Markus Sutter, Matt Melnicki, Ryan Leverenz, Emily Pawlowski (MSU)

Mike Guttman (U Wash)

Chris Petzold, Leanne Chan (JBEI)

Rich Celestre, Jun Feng (ALS)



# AGENDA



- Context: Why we care about the Orange Carotenoid Protein (OCP)
- What we've learned so far: hybrid methods lead to hypothesis for a photoswitch mechanism
- Future plans for elucidating the bigger system




# PHOTOPROTECTION



Photosynthetic organisms turn light into chemical energy through a complex interplay of proteins

The Phycobilisomes are the light-harvesting “antennae”




# PHOTOPROTECTION



But too much light can damage the organism



Cyanobacteria have evolved a unique way of dissipating excess energy using the Orange Carotenoid Protein (OCP)








# A KEY PLAYER: THE ORANGE CAROTENOID PROTEIN

- Activated in cyanobacteria in high light conditions
- Used to quench excess energy
- Contains a light-sensitive pigment carotenoid
- Inactive "orange" form has been crystallized
- Active "red" form has not been crystallized





# ORANGE TO RED CONVERSION VIA LIGHT









# PROTECTIONS OBSERVED USING FOOTPRINTING



# FOOTPRINTING DATA SUPPORTS DOMAIN SEPARATION




Gupta, Guttman et al, "Local and global structural drivers for the photoactivation of the orange carotenoid protein," *PNAS*, V112 No41, E5567, 2015.




# SMALL ANGLE SCATTERING (SAXS) AND HYDROGEN-DEUTERIUM EXCHANGE (HDX) DATA SUPPORT DOMAIN SEPARATION



## HDX



## SAXS





The model shows that in OCPR, the NTD and CTD are separated by ~ 16 Å







# CAROTENOID MOVEMENT WITHIN THE PROTEIN



Leverenz, Gupta et al,  
"Carotenoid translocation in  
the Orange Carotenoid  
Protein activates a  
photoprotective mechanism  
in cyanobacteria," *Science*  
V348, 6242, p1463, 2015.



# PROTECTION OF OCP WHEN FRP BINDS






# FOOTPRINTING FROM THE FRP “POINT OF VIEW”

Peptide	Mods	FRP / OCPO-FRP	FRP / OCPR-FRP (color on structure)
ELDGLLATVQAQA SQITQIDDLWK	Various residues	1	1.5 (light blue)
LHDFLSAK	Various residues	1.4	1.5 (light blue)
QSV <sup>II</sup> IFVFAQ <sup>LL</sup> KK	Predominant at II and LL	1.4	2.1 (dark blue)
EGLVQAEELTFLA ADK	Predominant at EE and F	1	1.2 (didn't plot)

Higher number means more protection upon complex formation



Color	Fold Protec.
Light Blue	1.5
Dark Blue	2.1







# TIME RESOLVED ORANGE/RED CONVERSION

Exposure time  
constant at: 0.3ms

R155,156



W41, Y44, M47

ORANGE TO RED



More modification

RED TO ORANGE




More modification






# PICTORAL VIEW OF ORANGE-RED CONVERSION

## Decrease in SA




## Increase in SA



$> 0.010 \text{ s}^{-1}$  (Fast)


$< 0.010$  to  $> 0.005 \text{ s}^{-1}$

$> 0.005 \text{ s}^{-1}$  (Slow)



## OCP<sup>O</sup> to OCP<sup>R</sup> Conversion

■ Increase in SA  
■ Decrease in SA







## WHAT'S NEXT?

- Repeat the kinetics experiments
- Investigate FRP light sensitivity
- Further investigate FRP-OCP interactions
- Investigate OCP-PB interactions

